

# 5-Fluorouracil cardiotoxicity: Molecular mechanisms and protective effects of simvastatin

Radwa Mohammed<sup>1</sup>, Nada Sallam, Hanan El Abhar

<sup>1</sup>Department of Pharmacology and Toxicology – Faculty of Pharmacy, Cairo University, Egypt  
radwa.nasser@pharma.cu.edu.eg

## Introduction

5-fluorouracil (5-FU) is a chemotherapeutic agent widely used in the treatment of different solid tumours, especially colorectal cancer. Its use is associated with rare but potentially serious cardiovascular toxicity. This study aims to investigate molecular mechanisms underlying the cardiovascular toxicity of 5-FU and the potential protective effects of simvastatin.

## Methods

Adult male albino Wistar rats were randomly divided into four groups (15-20/group).

- ▶ The first group received normal saline (i.p) once weekly for six successive weeks.
- ▶ In the second group rats received 5-FU (50 mg/kg; i.p) once weekly for six successive weeks (cardiotoxic group).
- ▶ Rats of the third group received simvastatin (15 mg/kg/day, p.o.) daily for eight successive weeks.
- ▶ Finally, rats of the fourth group received simvastatin daily a week before the first 5-FU injection, then concomitantly for six weeks, and continued alone for another week after the last dose of 5-FU.

◆ ECG recording was weekly carried out. Cardiac content of NADPH-oxidase, COX-2, NF-κB, p-eNOS and p-AKT in addition to aortic content of endothelin-1 and thromboxane-A2 were assessed by enzyme-linked immunosorbent assay. Protein expression of cardiac caspase-3 and Rho-kinase was evaluated by western blotting. Serum level of NT-proBNP and cardiac malondialdehyde (MDA) were also evaluated. Finally, histopathological evaluation of both cardiac and aortic tissues was carried out.

## Results

- ▶ 5-FU caused histopathological changes in both myocardial and aortic tissues.
- ▶ Myocardial ischemia and QTc prolongation were confirmed by ECG recording.
- ▶ 5-FU increased myocardial NADPH-oxidase and COX-2 content, leading to increased ROS production.
- ▶ Oxidative stress, inflammation and associated apoptosis in the heart were indicated by elevated MDA, NF-κB content and caspase-3 protein expression, respectively.
- ▶ Elevated aortic tissue content of endothelin-1 and thromboxane-A2, the two potent vasoconstrictors was observed.
- ▶ 5-FU significantly increased ROCK protein expression and p-AKT content, and suppressed p-eNOS level.
- ▶ Finally, elevated serum level of NT-proBNP was observed.
- ▶ Simvastatin was able to mitigate most of these abnormalities.

## Conclusion

- ▶ Direct myocardial injury and ischemia caused by endothelial dysfunction and activation of Rho/ROCK pathway together with suppressed endothelial NOS activity are potential mechanisms of 5-FU cardiovascular toxicity.
- ▶ Inhibition of ROCK activity by simvastatin, a drug with potent antioxidant and pleiotropic properties, mitigates the cardiovascular toxicity of 5-FU.
- ▶ Elevated serum level of NT-proBNP associated with 5-FU treatment may be used as a marker of cardiotoxicity, but further studies are needed to confirm or negate this.
- ▶ Elevated tissue levels of endothelin-1 and thromboxane-A2 may also account for the vasoconstrictor and ischemic response of the body towards 5-FU therapy.





Figure 1: The proposed molecular mechanisms that can account for 5-FU cardiotoxicity in the present study.



1. Vincent DT et al., The role of antioxidants in the era of cardio oncology. *Cancer Chemother Pharmacol* (2007) 1157-1168.
2. Albini A et al., Cardiotoxicity of anticancer drugs: the need for cardio-oncology and cardio-oncological prevention. *J Natl Cancer Inst* (2009) 14-25.
3. Jensen SA et al., Fluorouracil induces myocardial ischemia with increases of plasma brain natriuretic peptide and lactic acid but without dysfunction of left ventricle. *J Clin Oncol* (2010) 5280-5286.
4. Shimokawa H RhoA/Rho-Kinase in the Cardiovascular System. *Circ Res* 2016 (118) 352-366.