

Survivin and Caspase-3 as a diagnostic and predictive biomarkers of recurrence for urinary bladder carcinoma after TURBT

Vivian GD Rouston¹, Amal AA Shaaban², Dina M Abd Allah² and Ahmed F Kotb²

¹*Ministry of Health, Egypt*

²*Alexandria University, Egypt*

Abstract

Background: Poor sensitivity of cytology and invasiveness of urethroscopy have generated interest in non-invasive tools to monitor for recurrence. Caspase-3 and survivin have central role in regulation of apoptosis. Survivin can aid early diagnosis, determine prognosis in multiple cancer types and predict response to anti-cancer therapies. Its combination with other biomarkers as caspase-3 enhance prognostication and prediction of treatment response in UBC.

Methods: Immunohistochemical expression of survivin and caspase-3 were assessed in 44 Egyptian consecutive patients with UBC and 7 cystoscopic biopsies of cystitis as control reactive benign urothelium. Relationships between their expression, clinicopathological characteristics, diagnostic and prognostic performance were statistically analyzed.

Findings: No survivin immunoreactivity was identified in non-neoplastic bladder tissue. Expression of survivin and caspase-3 was altered in 42(95.5%) and 10(22.7%) cases, respectively. There was statistically significant moderate positive correlation between survivin and caspase-3 expression among whole studied cases ($p=.006$). Expression of either survivin or caspase-3 protein individually significantly differ ($p=0.000$) in cancer status from control cases. Survivin was an independent predictor of UBC in multivariable analyses. Diagnostic accuracy of survivin alone was significantly better than caspase-3 alone (sensitivity 81.82% Vs 68.18%, $p=.027$). Addition of survivin immunoreactivity to model including caspase-3 expression improved diagnostic accuracy with a sensitivity of 93.18%. Addition of gender to the previous model improved more diagnostic accuracy with sensitivity of 100%.

Interpretation: Survivin alone is very promising marker and reliable indicator in UBC. Survivin and caspase-3 antigens have a cooperative effect on bladder cancer, their simultaneous evaluation augments diagnostic sensitivity.

Abbreviations: UBC, Urinary Bladder Cancer or Carcinoma; TURBT, Trans Urethral Resection of Bladder Tumor




Figure 1: IHC expression of survivin. (a) Normal survivin immunostaining 7% in a chronic non specific cystitis (X100). (b) Altered survivin immunostaining 90% in a low grade papillary UC, showing diffuse nuclear predominance staining and moderate cytoplasmic staining pattern (X100). (c) Altered survivin immunostaining 70% in a high grade papillary UC, showing diffuse cytoplasmic immunopositivity (X200). (d) Altered survivin expression 70% in a low grade non-papillary UC, showing focal cytoplasmic and nuclear immunostaining (X100). (e & f) Altered survivin immunostaining 80% in a high grade non papillary UC, showing diffuse cytoplasmic staining (X100 & X400).




Figure 2: Caspase-3 immunoreactivity. (a) Caspase-3 normal "15%" cytoplasmic immunostaining in chronic cystitis with von brunn nests (X400). (b) Altered "70%" Caspase-3 immunostaining in a low grade papillary UC, showing diffuse cytoplasmic staining (X200). (c) Caspase-3 expression 70% in a low grade non papillary UC, showing nuclear and cytoplasmic staining (X100). (d) Caspase-3 expression 90% in a high grade papillary UC, showing diffuse cytoplasmic staining (X400). (e) Caspase-3 immunostaining 70% in a high grade non papillary UC, showing diffuse cytoplasmic staining pattern (X400).




Figure 3: ROC curve comparing the performance of caspase-3 expression alone, survivin expression alone, Model one (caspase-3 and survivin expression) and Model two (caspase-3 immunoreactivity, survivin expression and gender) in diagnosis of urinary bladder cancer among studied sample.

Biography

Vivian GD Rouston has obtained her MBChB degree from Faculty of Medicine, Alexandria University. She has pursued her MSc degree in Anatomic Pathology from Faculty of Medicine Alexandria University. She is currently working as a Histopathology Specialist in a general hospital of the Egyptian Ministry of Health.

v_dabous@hotmail.com