

# Vanadium and chromium mediated impairments in the immunological reactivity of rats with aseptic inflammation

Aliya N. Tokusheva, Department of Pathophysiology, Asfendiyarov Kazakh National Medical University, Almaty, Kazakhstan

Marina K. Balabekova, Candidate of Medical Sciences, Department of Pathophysiology, Asfendiyarov Kazakh National Medical University, Almaty, Kazakhstan

Yekaterina O. Ostapchuk, PhD, M.A.Aitkhozhin's Institute of Molecular Biology and Biochemistry, Laboratory of Molecular Immunology and Immunobiotechnology, Almaty, Kazakhstan

Nikolay N. Belyaev, Doctor of Biological Sciences, Professor, M.A.Aitkhozhin's Institute of Molecular Biology and Biochemistry, Laboratory of Molecular Immunology and Immunobiotechnology, Almaty, Kazakhstan

Rustam R. Tukhvatshin, Doctor of Medical Sciences, Department of Pathophysiology, Kyrgyz State Medical Academy named after I.K. Akhunbaev, Bishkek, Kyrgyzstan

## Abstract

Chronic inflammation is a consequence of the immune system dysfunction. Such dysregulation of the immune response may induce chronic diseases, such as autoimmune diseases, diabetes, and malignant transformation of cells. Xenobiotics, including vanadium and chromium, were shown to induce inflammatory changes leading to chronic inflammation. The purpose of this research was to study the process of aseptic inflammation accompanied with intoxication with salts of heavy metals. In our study, sexually mature rats were administered aseptic inflammation (AI) alone or ammonium vanadate and potassium dichromate (AV/PD) at a dose of 5 mg/kg of BW for two weeks and after that aseptic inflammation was modeled. Lymphatic organs were studied on day 1, 7, and 14 after the onset of aseptic inflammation. Administration of AV/PD and AI resulted in structural changes in lymphatic organs and anemia observed throughout the experiment. We observed a decrease in the cellularity of the bone marrow and thymus, ratio of thymic cortex to medulla, and dystrophic changes in thymic cells and their scarcity. Also, we detected increased levels of anti-inflammatory cytokines in serum: IL-10 on day 1 and TGF- $\beta$  on day 7 and 14 after the beginning of the experiment in the group AI+AV/PD comparing to intact rats and AI group. Phenotypical analysis demonstrated that by the end of experiment freshly obtained splenocytes of AI and AI+AV/PD rats contained increased percentage of His48<sup>+</sup>CD11b/c<sup>+</sup> and His48<sup>high</sup>CD11b/c<sup>+</sup> cells, and decreased number of induced CD3<sup>+</sup>CD4<sup>+</sup>IFN $\gamma$ <sup>+</sup>, CD3<sup>+</sup>CD4<sup>+</sup>IL-4<sup>+</sup> and CD3<sup>+</sup>CD8<sup>+</sup> cells, comparing to control animals. Interestingly, during the next period of the experiment, we observed significant decrease of induced CD3<sup>+</sup>CD4<sup>+</sup>IFN $\gamma$ <sup>+</sup>, CD3<sup>+</sup>CD4<sup>+</sup>IL-4<sup>+</sup> and CD3<sup>+</sup>CD8<sup>+</sup> cells in the AI+AV/PD group comparing to AI group.

Thus, it is possible that an elevated sera IL-10 and TGF- $\beta$ 1 observed in mice with chronic inflammatory processes administered with AV/PD result of abundant accumulation of His48<sup>+</sup>CD11b/c<sup>+</sup> and His48<sup>high</sup>CD11b/c<sup>+</sup> cells myeloid cells in the periphery, and, in turn, it could participate in the maintaining of the immunosuppressive environment that supports persistence of chronic inflammatory conditions. Therefore, intoxication with vanadium and chromium salts may support immunosuppressive environment, contributing to chronic inflammation development.

## Image




Figure. Splenocytes, obtained from control rats, AI, and AI+AV/PD groups on day 1, 7, and 14 after the beginning of the experiment, were labeled with anti-His48 and anti-CD11b/c, after that cells were analyzed by flow cytometry. Each group included 6 rats. Representative flow cytometry data for each group are depicted.

## Recent Publications (minimum 5)

1. Chandra AK, Ghosh R, Chatterjee A, Sarkar M (2007) Effects of vanadate on male rat reproductive tract histology, oxidative stress markers and androgenic enzyme activities. *J. Inorganic Biochem* 101(6):944–956.
2. Rose-John S (2012) IL-6 Trans-Signaling via the Soluble IL-6 Receptor: Importance for the Pro-Inflammatory Activities of IL-6. *Int J Biol Sci* 8(9):1237–1247.
3. deSeny D, Cobraiville G, Leprince P *et al.* (2016) Biomarkers of inflammation and innate immunity in atrophic nonunion fracture. *J Transl Med* 14:258-268.
4. Schwartz C, O'Grady K, Lavelle EC, Fallon PG (2016) Interleukin 33: an innate alarm for adaptive responses beyond Th2 immunity-emerging roles in obesity, intestinal inflammation, and cancer. *Eur J Immunol* 46 (5):1091-1100.
5. Huang Y, Chen Z (2016) Inflammatory bowel disease related innate immunity and adaptive immunity. *Am J Transl Res* 15 (6):2490-2497.



## Biography

Aliya Tokusheva is a PhD student at the Asfendiyarov Kazakh National Medical University. She studies molecular mechanisms of epigenetic regulation of tissue-specific expression of genes, revealing the effect of heavy metal compounds on the variability of the genome of immunogenesis organs and the mutual regulatory influence of

important links of these mechanisms on the course of the inflammatory process.

Email: [aliyatokusheva@mail.ru](mailto:aliyatokusheva@mail.ru)

---