



PREVALENCE OF HIGH TITRE ALPHA AND BETA HAEMOLYSINS AMONG BLOOD DONORS IN SOKOTO, NORTH WESTERN NIGERIA

PRESENTER:

PROF. EMMANUEL K. UKO

Prof. of Haematology and Blood Transfusion Science,
College of Medicine, Department of Medical Laboratory
Science, University of Calabar, Nigeria

Email: emmanuelkuko@gmail.com

BACKGROUND

Allogenic blood has never been more in demand than it is today particularly in developing countries. One of the biggest challenges to blood safety particularly in developing countries is accessing safe and adequate quantities of blood and blood products. One strategy geared towards the optimum utilization of scarce donor units is use of blood and blood products against ABO blood group barrier.

The aim of this study area was to determine the prevalence of alpha and beta- haemolysins among blood groups A, B and O donors in Sokoto, North Western, Nigeria

METHODS

This prospective case study was carried out at the service Laboratory of the Faculty of Medical Laboratory science in Usmanu Danfodiyo University Sokoto, North Western Nigeria. Sera from 140 blood donors were screened for alpha- (anti-A) and beta- (anti-B) haemolysins using the standard tube technique at 37°C for 1 hour. Titre of ≥ 1.64 was regarded as positive for high titre haemolysin.

RESULT

One hundred and forty blood males donors aged 18 – 50 years and mean age 30 ± 8.4 years were screened for alpha- (anti-A) and beta- (anti-B) haemolysins.

Study population included 60 blood group O donors (42.9%) and 40 blood group A (26.7%) and 40 blood group B (26.7%) donors. Out of the 140 blood donors screened, 14(10.0%) were positive for high titre haemolysin (titre ≥ 64) while 126 (90%) were negative (titre < 64).

The prevalence of haemolysin was compared based on blood group of donors. The prevalence of high titre haemolysin was significantly higher among blood group O donors (18.3%) compared to 5% and 2,5% respectively for group B and group A donors respectively (= 0.001).

Out of the 14 donors positive for haemolysin, 7 (50%) were positive for Alpha haemolysin, 4(28.5%) for beta while 3(21.4) had both alpha and beta haemolysin. There was no age dependent differences in the prevalence of high titre haemolysins among the blood donor tested.

Table 1: Distribution of ABO blood group and Prevalence of haemolysin among the 140 blood donors studied

ABO Blood Group	No. of Subjects	% Distribution of the Groups	Prevalence of Hemolysis
O	60	42.9%	18.3%
A	40	28.6%	2.5%
B	40	28.6%	5.0%

Table 2: Prevalence of High Titre Haemolysin among the 140 blood donors Screened

No. of Donors	Titre (%)
14	> 64 (10)
126	<64 (90)

Table 3: Distribution of α , β and $\alpha\beta$ Hemolysis in the Donor with high Titre

Haemolysin Type	No. of Positivity
α	7
β	4
$\alpha\beta$	3

CONCLUSION

This study has shown that the prevalence of high titre alpha and beta haemolysin is high particularly among blood group O donors in Sokoto, Nigeria. We recommend that all group O, A and B blood whose blood is intended for transfusion and ABO blood group barrier be screened routinely for high titre alpha and beta haemolysins.

All blood group O red cells unit intended for use as universal donor units must be screened and found negative for high titre alpha and beta haemolysins.

Also all fresh frozen plasma (FFP) and platelet concentrate intended for use against ABO blood group barrier should be tested and found negative for high titre anti-A and B haemolysin.

Thank you!