



EFFECTIVENESS OF SCALING AND ROOT PLANNING TREATMENT FOR SEVERE CHRONIC PERIODONTITIS ALONG WITH SIMULTANEOUS USE OF THREE DIFFERENT ANTIBIOTICS AND IRRIGATION SOLUTIONS

Visar Bunjaku DDS^{1*}, Aneta Atanasovska – Stojanovska DDS PhD², Mirjana Popovska DDS PhD², Shefqet Mrasori DDS PhD³, Metush Disha DDS PhD⁴, Fatmir Dragidella DDS PhD⁴, Zana Sllamniku DDS PhD⁴, Arben Murtezani DDS⁵

Background

- ▣ Purpose of this prospective study was to evaluate the effectiveness of systemic administration of three different antibiotics and irrigation solutions following scaling and root planning (SRP) for the treatment of severe chronic periodontitis.

Materials and Methods

- ▣ 114 patients (age group 29 – 57) were selected and randomly assigned to scaling and root planning with at least seven teeth with probing depth ≥ 5 mm and bleeding on probing (BoP).

Materials and Methods

- ▣ Patients are divided into three groups;
- ▣ Group A received SRP + irrigation using 2% peroxide solution followed by systemic use of amoxicillin 1000 mg twice a day for 10 days.
- ▣ Group B received SRP + irrigation with 0.2% chlorexidine solution followed by systemic use of metronidazole 400 mg twice a day for 10 days .

Materials and Methods

- ▣ Group C received SRP + irrigation with 3% Povidone-Iodine solution followed by systemic use of Azythromycine 500 mg once a day for 5 days.
- ▣ Gain in clinical attachment level (CAL), probing depth (PD), bleeding on probing (BoP) and visible plaque index were selected as outcome variables.

Materials and Methods

- ▣ Weekly, proficient supragingival plaque removal was performed.
- ▣ Following all therapies, a reduction in full mouth mean clinical parameters was observed at baseline and after 60 days by the same examiner.



"You're teeth look great. But it looks like we need to extract your tongue."

Results

- ▣ All three groups have improvements in clinical parameters compared to baseline ($p < 0.05$).
- ▣ The mean probing depth reductions were:
Group A = 0.70 mm, Group B = 0.85 mm and
Group C = 1.10 mm.

Results

- ▣ The probing depth reduction (PD) and gain in clinical attachment level (CAL) was considerably higher in the SRP, Povidone - Iodine irrigation plus Azythromycine group, compare to two other groups ($p=0.003$).

Results

- ▣ As far as bleeding on probing (BoP) and visible plaque index there was no significant difference between the groups ($p < 0.05$).

Conclusion

- ▣ Results of this short term study demonstrated significant advantages of systemic use of Azytromycine after SRP and irrigation with 3% Povidone - Iodine solutions for the treatment of severe chronic periodontitis.

Thank You !

