

Abstract (600 word limits)

Non-endoscopic minimally invasive evacuation of intracerebral haematoma

^{1,2}Tomaz Velnar

¹Department of neurosurgery, University medical centre Ljubljana, Slovenia

²AMEU, Alma Mater Europaea university Maribor, Slovenija

Spontaneous intracerebral haemorrhage has a high disability and mortality rate. In cases, when surgery is needed, minimally invasive approach is recommended.

A 59-year old patient was admitted due to progressive left sided arm and leg weakness. The neurological status started to deteriorate quickly. A computed tomography (CT) of the head revealed an ICH of 7cm in diameter with haematocephalus and cerebral oedema. The CT angiography was negative, classifying the haematoma as a primary one. Coagulation and aggregation values were deranged as a result of liver failure. The international normalised ratio (INR) and prothrombine time (PT) were lowered to 1.56 and 0.47, respectively. The platelet count was 33 and the platelet function tests were completely disturbed. Injections of fresh frozen plasma, recombinant coagulation factor VIIa, protrombin complex, vitamin K and platelet plasma were applied. As a result of extensive intracerebral bleeding and consciousness decline, surgery was recommended despite unfavourable laboratory results. A minimally invasive approach was chosen for the ICH removal. A burr hole of 1cm in diameter was made in the right temporal area. Under the microscope, the liquefied blood was evacuated with aspirator and bipolar. The ICP values remained normal during the course of treatment. The control CT scan showed successfully evacuated haematoma and normal width of the ventricles. The sedation was gradually discontinued after a week. The patient was awake with persistent left sided haemiplegia.

In case of patient with numerous risk factors and imminent operation, minimally invasive surgery for intracerebral haematoma is warranted.

Biography (200 word limit)

Tomaz Velnar, MD, PhD is a neurosurgeon and assistant professor at Ljubljana medical centre. He is also active in research, cooperating regularly with the other two authors. They have started a multicentre study of vitamin D deficiency among older people

References

1. Bilezikian, J. P., Cusano, N. E., Khan, A. A., Liu, J., Marcocci, C., & Bandeira, F. (2016, May 19). [Primary hyperparathyroidism.](#)

2. Doshi, M., Lahoti, A., Danesh, F. R., Batuman, V., & Sanders, P. W. (2016, December 07). [Paraprotein-Related Kidney Disease: Kidney Injury from Paraproteins—What Determines the Site of Injury?](#)

3. Gerecke, C., Fuhrmann, S., Striffler, S., Schmidt-Hieber, M., Einsele, H., & Knop, S.

(2016, July). [The Diagnosis and Treatment of Multiple Myeloma](#). Rajkumar, S. V. (2016, July). Multiple Myeloma: 2016 update on Diagnosis, Risk- stratification and Management.

Organization / University Logo

